

Instructions: You should have with you several number two pencils, an eraser, your 3" x 5" note card, a calculator, and your University ID Card. If you have notes with you, place them in a sealed backpack and place the backpack OUT OF SIGHT or place the notes directly on the table at the front of the room.

Fill in the front page of the Scantron answer sheet with your class section number (see below), last name, first name, middle initial, and student identification number. Leave the test form number blank.

Section 001 (MWF 8am with Dr. Nafshun)
Section 003 (MWF 11am with Dr. Watson)
Section 005 (MWF 2pm with Dr. Grajczyk)


Section 002 (MWF 9am with Dr. Nafshun)
Section 004 (MWF 1pm with Dr. Burrows)

This exam consists of 32 multiple-choice questions; each has 5 points attached. When you finish this exam, proceed to the proctor. Show your OSU ID Card and submit your completed Scantron form. You may take your notecard and this exam packet with you.

$K_a[\text{CH}_3\text{COOH (aq)}] = 1.80 \times 10^{-5}$ (acetic acid)	$K_a[\text{C}_6\text{H}_5\text{COOH (aq)}] = 6.30 \times 10^{-5}$ (benzoic acid)
$K_a[\text{CH}_2\text{ClCOOH (aq)}] = 1.40 \times 10^{-3}$ (chloroacetic acid)	$K_b[\text{NH}_3 \text{ (aq)}] = 1.80 \times 10^{-5}$ (ammonia)
$K_a[\text{HClO (aq)}] = 2.90 \times 10^{-8}$ (hypochlorous acid)	$K_a[\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{COOH (aq)}] = 1.45 \times 10^{-5}$ (pentanoic acid)
$K_a[\text{HF (aq)}] = 6.30 \times 10^{-4}$ (hydrofluoric acid)	$K_b [\text{CH}_3\text{NH}_2] = 3.70 \times 10^{-4}$ (methylamine)
$K_a[\text{HCOOH (aq)}] = 1.80 \times 10^{-4}$ (formic acid)	$K_{sp} [\text{Fe(OH)}_2] = 4.87 \times 10^{-17}$
$K_{sp} [\text{PbF}_2] = 3.6 \times 10^{-8}$	$K_{sp} [\text{MgF}_2] = 3.7 \times 10^{-8}$
$K_{sp} [\text{Cd(OH)}_2] = 7.2 \times 10^{-15}$	$K_{sp} [\text{PbI}_2] = 1.4 \times 10^{-8}$
$K_{sp} [\text{CaSO}_4] = 2.4 \times 10^{-5}$	$K_{sp} [\text{CaC}_2\text{O}_4] = 2.3 \times 10^{-9}$
$K_{sp} [\text{CuCl}] = 1.0 \times 10^{-6}$	$K_{sp} [\text{AgCl}] = 1.77 \times 10^{-10}$
$K_{sp} [\text{Ag}_2\text{CO}_3] = 8.1 \times 10^{-12}$	

First-Order		
$\ln [A] = -kt + \ln [A]_0$	$t_{1/2} = \frac{\ln (2)}{k}$	$\ln \frac{A}{A_0} = -kt$

Water	$\text{H}-\overset{\cdot\cdot}{\underset{\cdot\cdot}{\text{O}}}-\text{H}$	Note: No Charge
Ammonia	$\begin{array}{c} \text{H}-\overset{\cdot\cdot}{\underset{\cdot\cdot}{\text{N}}}-\text{H} \\ \\ \text{H} \end{array}$	Note: No Charge
Chloride ion	$[\overset{\cdot\cdot}{\underset{\cdot\cdot}{\text{Cl}}}]^{-}$	Note: Minus one charge
Carbon monoxide	$:\text{C}\equiv\text{O}:$	Note: No Charge
Cyanide ion	$[\text{C}\equiv\text{N}]^{-}$	Note: Minus one charge
Thiocyanate ion	$[\overset{\cdot\cdot}{\underset{\cdot\cdot}{\text{S}}}=\text{C}=\overset{\cdot\cdot}{\underset{\cdot\cdot}{\text{N}}}]^{-}$	Note: Minus one charge
Oxalate ion (ox)	$[\overset{\cdot\cdot}{\underset{\cdot\cdot}{\text{O}}}=\overset{\cdot\cdot}{\underset{\cdot\cdot}{\text{C}}}-\overset{\cdot\cdot}{\underset{\cdot\cdot}{\text{C}}}=\overset{\cdot\cdot}{\underset{\cdot\cdot}{\text{O}}}]^{2-}$	Note: Minus two charge
Ethylenediamine (en)	$\begin{array}{c} \text{H} & \text{H} \\ & \\ \text{H}-\overset{\cdot\cdot}{\underset{\cdot\cdot}{\text{N}}}-\text{C}-\text{C}-\overset{\cdot\cdot}{\underset{\cdot\cdot}{\text{N}}}-\text{H} \\ & \\ \text{H} & \text{H} \end{array}$	Note: No Charge
Ethylenediaminetetraacetate (EDTA)	$[\overset{\cdot\cdot}{\underset{\cdot\cdot}{\text{O}}}-\overset{\cdot\cdot}{\underset{\cdot\cdot}{\text{C}}}\overset{\cdot\cdot}{\underset{\cdot\cdot}{\text{O}}}-\text{CH}_2-\overset{\cdot\cdot}{\underset{\cdot\cdot}{\text{N}}}-\text{CH}_2-\text{CH}_2-\overset{\cdot\cdot}{\underset{\cdot\cdot}{\text{N}}}-\text{CH}_2-\overset{\cdot\cdot}{\underset{\cdot\cdot}{\text{C}}}\overset{\cdot\cdot}{\underset{\cdot\cdot}{\text{O}}}-\overset{\cdot\cdot}{\underset{\cdot\cdot}{\text{O}}}]^{4-}$	Note: Minus four charge



Spectrochemical series: $\text{CN}^- > \text{NO}_2^- > \text{en} > \text{NH}_3 > \text{NCS}^- > \text{H}_2\text{O} > \text{OH}^- > \text{F}^- > \text{Cl}^-$

$$R = 8.314 \text{ J/mol}\cdot\text{K}$$

$$N_A = 6.022 \times 10^{23}$$

$$h = 6.626 \times 10^{-34} \text{ J}\cdot\text{s/photon}$$

$$c = 3.00 \times 10^8 \text{ m/s}$$

$$K = ^\circ\text{C} + 273.15$$

$$E = hc/\lambda$$

The Standard Molar Entropies of Selected Substances at 298.15 K (25°C)

Compound	$S_m^\circ / \text{J K}^{-1} \text{mol}^{-1}$	Compound	$S_m^\circ / \text{J K}^{-1} \text{mol}^{-1}$
Solids		Diatomic Gases	
C (diamond)	2.377	H ₂	130.7
C (graphite)	5.74	D ₂	145.0
Si	18.8	HCl	186.9
Ge	31.1	HBr	198.7
Sn (gray)	44.1	HI	206.6
Pb	64.8	N ₂	191.6
Li	29.1	O ₂	205.1
Na	51.2	F ₂	202.8
K	64.2	Cl ₂	223.1
Rb	69.5	Br ₂	245.6
Cs	85.2	I ₂	260.7
NaF	51.5	CO	197.7
MgO	26.9	Triatomic Gases	
AlN	20.2	H ₂ O	188.8
NaCl	72.1	NO ₂	240.1
KCl	82.6	H ₂ S	205.8
Mg	32.7	CO ₂	213.7
Ag	42.6	SO ₂	248.2
I ₂	116.1	N ₂ O	219.9
MgH ₂	31.1	O ₃	238.9
AgN ₃	99.2	Polyatomic Gases (> 3)	
Liquids		CH ₄	186.3
Hg	76.0	C ₂ H ₆	229.6
Br ₂	152.2	C ₃ H ₈	269.9
H ₂ O	69.9	C ₄ H ₁₀	310.2
H ₂ O ₂	109.6	C ₅ H ₁₂	348.9
CH ₃ OH	126.8	C ₂ H ₄	219.6
C ₂ H ₅ OH	160.7	N ₂ O ₄	304.3
C ₆ H ₆	172.8	B ₂ H ₆	232.0
BCl ₃	206.3	BF ₃	254.0
Monatomic Gases		NH ₃	192.5
He	126.0		
Ne	146.2		
Ar	154.8		
Kr	164.0		
Xe	169.6		

$$F = 96,485 \text{ C/mole } e^-$$

$$R = 8.314 \text{ J/mol}\cdot\text{K}$$

$$N_A = 6.022 \times 10^{23}$$

$$S = k \ln W$$

$$h = 6.626 \times 10^{-34} \text{ J}\cdot\text{s/photon}$$

$$\Delta G^\circ_{\text{rxn}} = -RT \ln K$$

$$\ln K = -\frac{\Delta H^\circ_{\text{rxn}}}{R} \left(\frac{1}{T}\right) + \frac{\Delta S^\circ_{\text{rxn}}}{R}$$

$$\Delta G_{\text{rxn}} = \Delta G^\circ_{\text{rxn}} + RT \ln Q$$

$$c = 3.00 \times 10^8 \text{ m/s}$$

$$E_{\text{cell}} = E^\circ_{\text{cell}} - \frac{RT}{nF} \ln Q$$

$$\Delta G^\circ = -nF E^\circ_{\text{cell}}$$

$$\Delta G = \Delta H - T\Delta S$$

$$k = 1.381 \times 10^{-23} \text{ J/K}$$

$$K = ^\circ\text{C} + 273.15$$

$$E = hc/\lambda$$

$$t_{1/2} [^{14}\text{C}] = 5730 \text{ y}$$

Reduction Half-Reaction	E°, volt
Acidic Solution	
$F_2(g) + 2 e^- \rightarrow 2F^-(aq)$	+2.866
$O_3(g) + 2 H^+(aq) + 2 e^- \rightarrow O_2(g) + H_2O(l)$	+2.075
$S_2O_8^{2-}(aq) + 2 e^- \rightarrow 2 SO_4^{2-}(aq)$	+2.01
$H_2O_2(aq) + 2H^+(aq) + 2 e^- \rightarrow 2 H_2O(l)$	+1.763
$MnO_4^-(aq) + 8H^+(aq) + 5 e^- \rightarrow Mn^{2+}(aq) + 4 H_2O(l)$	+1.51
$PbO_2(s) + 4H^+(aq) + 2 e^- \rightarrow Pb^{2+}(aq) + 2 H_2O(l)$	+1.455
$Cl_2(g) + 2 e^- \rightarrow 2 Cl^-(aq)$	+1.358
$Cr_2O_7^{2-}(aq) + 14 H^+(aq) + 6 e^- \rightarrow 2 Cr^{3+}(aq) + 7 H_2O(l)$	+1.33
$MnO_2(s) + 4H^+(aq) + 2 e^- \rightarrow Mn^{2+}(aq) + 2 H_2O(l)$	+1.23
$O_2(g) + 4H^+(aq) + 4 e^- \rightarrow 2 H_2O(l)$	+1.229
$2 IO_3^-(aq) + 12H^+(aq) + 10 e^- \rightarrow I_2(s) + 6 H_2O(l)$	+1.20
$Br_2(l) + 2 e^- \rightarrow 2 Br^-(aq)$	+1.065
$NO_3^-(aq) + 4H^+(aq) + 3 e^- \rightarrow NO(g) + 2 H_2O(l)$	+0.956
$Ag^+(aq) + e^- \rightarrow Ag(s)$	+0.800
$Fe^{3+}(aq) + e^- \rightarrow Fe^{2+}(aq)$	+0.771
$O_2(g) + 2H^+(aq) + 2 e^- \rightarrow H_2O_2(aq)$	+0.695
$I_2(s) + 2 e^- \rightarrow 2 I^-(aq)$	+0.535
$I_3^-(aq) + 2 e^- \rightarrow 3 I^-(aq)$	+0.530
$Cu^{2+}(aq) + 2 e^- \rightarrow Cu(s)$	+0.340
$SO_4^{2-}(aq) + 4H^+(aq) + 2 e^- \rightarrow 2 H_2O(l) + SO_2(g)$	+0.17
$Sn^{4+}(aq) + 2 e^- \rightarrow Sn^{2+}(aq)$	+0.154
$S(s) + 2H^+(aq) + 2 e^- \rightarrow H_2S(g)$	+0.14
$2H^+(aq) + 2 e^- \rightarrow H_2(g)$	0
$Pb^{2+}(aq) + 2 e^- \rightarrow Pb(s)$	-0.125
$Sn^{2+}(aq) + 2 e^- \rightarrow Sn(s)$	-0.137
$Co^{2+}(aq) + 2 e^- \rightarrow Co(s)$	-0.277
$Cr^{3+}(aq) + e^- \rightarrow Cr^{2+}(aq)$	-0.410
$Fe^{2+}(aq) + 2 e^- \rightarrow Fe(s)$	-0.440
$Zn^{2+}(aq) + 2 e^- \rightarrow Zn(s)$	-0.763
$Al^{3+}(aq) + 3 e^- \rightarrow Al(s)$	-1.676
$Mg^{2+}(aq) + 2 e^- \rightarrow Mg(s)$	-2.356
$Na^+(aq) + e^- \rightarrow Na(s)$	-2.713
$Ca^{2+}(aq) + 2 e^- \rightarrow Ca(s)$	-2.84
$K^+(aq) + e^- \rightarrow K(s)$	-2.924
$Li^+(aq) + e^- \rightarrow Li(s)$	-3.040
Basic Solution	
$O_3(g) + H_2O(l) + 2 e^- \rightarrow O_2(g) + 2 OH^-(aq)$	+1.246
$OCl^-(g) + H_2O(l) + 2 e^- \rightarrow Cl^-(aq) + 2 OH^-(aq)$	+0.890
$O_2(g) + 2 H_2O(l) + 4 e^- \rightarrow 4 OH^-(aq)$	+0.401
$2 H_2O(l) + 2 e^- \rightarrow H_2(g) + 2 OH^-(aq)$	-0.828

The Periodic Table of the Elements

1 H Hydrogen 1.00794																	2 He Helium 4.003
3 Li Lithium 6.941	4 Be Beryllium 9.012182											5 B Boron 10.811	6 C Carbon 12.0107	7 N Nitrogen 14.00674	8 O Oxygen 15.9994	9 F Fluorine 18.9984032	10 Ne Neon 20.1797
11 Na Sodium 22.989770	12 Mg Magnesium 24.3050											13 Al Aluminum 26.981538	14 Si Silicon 28.0855	15 P Phosphorus 30.973761	16 S Sulfur 32.066	17 Cl Chlorine 35.4527	18 Ar Argon 39.948
19 K Potassium 39.0983	20 Ca Calcium 40.078	21 Sc Scandium 44.955910	22 Ti Titanium 47.867	23 V Vanadium 50.9415	24 Cr Chromium 51.9961	25 Mn Manganese 54.938049	26 Fe Iron 55.845	27 Co Cobalt 58.933200	28 Ni Nickel 58.6934	29 Cu Copper 63.546	30 Zn Zinc 65.39	31 Ga Gallium 69.723	32 Ge Germanium 72.61	33 As Arsenic 74.92160	34 Se Selenium 78.96	35 Br Bromine 79.904	36 Kr Krypton 83.80
37 Rb Rubidium 85.4678	38 Sr Strontium 87.62	39 Y Yttrium 88.90585	40 Zr Zirconium 91.224	41 Nb Niobium 92.90638	42 Mo Molybdenum 95.94	43 Tc Technetium (98)	44 Ru Ruthenium 101.07	45 Rh Rhodium 102.90550	46 Pd Palladium 106.42	47 Ag Silver 107.8682	48 Cd Cadmium 112.411	49 In Indium 114.818	50 Sn Tin 118.710	51 Sb Antimony 121.760	52 Te Tellurium 127.60	53 I Iodine 126.90447	54 Xe Xenon 131.29
55 Cs Cesium 132.90545	56 Ba Barium 137.327	57 La Lanthanum 138.9055	72 Hf Hafnium 178.49	73 Ta Tantalum 180.9479	74 W Tungsten 183.84	75 Re Rhenium 186.207	76 Os Osmium 190.23	77 Ir Iridium 192.217	78 Pt Platinum 195.078	79 Au Gold 196.96655	80 Hg Mercury 200.59	81 Tl Thallium 204.3833	82 Pb Lead 207.2	83 Bi Bismuth 208.98038	84 Po Polonium (209)	85 At Astatine (210)	86 Rn Radon (222)
87 Fr Francium (223)	88 Ra Radium (226)	89 Ac Actinium (227)	104 Rf Rutherfordium (261)	105 Db Dubnium (262)	106 Sg Seaborgium (263)	107 Bh Bohrium (262)	108 Hs Hassium (265)	109 Mt Meitnerium (266)	110 Ds Darmstadtium (269)	111 Rg Roentgenium (272)	112 Cn Copernicium (277)	113 Nh Nihonium	114 Fl Flerovium				

58 Ce Cerium 140.116	59 Pr Praseodymium 140.90768	60 Nd Neodymium 144.24	61 Pm Promethium (145)	62 Sm Samarium 150.36	63 Eu Europium 151.964	64 Gd Gadolinium 157.25	65 Tb Terbium 158.92534	66 Dy Dysprosium 162.50	67 Ho Holmium 164.93032	68 Er Erbium 167.26	69 Tm Thulium 168.93421	70 Yb Ytterbium 173.04	71 Lu Lutetium 174.967
90 Th Thorium 232.0381	91 Pa Protactinium 231.03588	92 U Uranium 238.0289	93 Np Neptunium (237)	94 Pu Plutonium (244)	95 Am Americium (243)	96 Cm Curium (247)	97 Bk Berkelium (247)	98 Cf Californium (251)	99 Es Einsteinium (252)	100 Fm Fermium (257)	101 Md Mendelevium (258)	102 No Nobelium (259)	103 Lr Lawrencium (262)


1. A student titrates 2.884 g of an unknown monoprotic acid to the equivalence point with 62.55 mL of 0.2447 M NaOH (aq). What is the molar mass of the acid?

- (A) 0.005307 g/mol
(B) 53.07 g/mol
(C) 530.7 g/mol
(D) 73.72 g/mol
(E) 188.4 g/mol

$$M_{\text{NaOH}} V_{\text{NaOH}} = \frac{\text{grams Acid}}{\text{MM Acid}}$$
$$(0.2447 \frac{\text{mol}}{\text{L}})(0.06255 \text{ L}) = \frac{2.884 \text{ g}}{\text{MM}}$$
$$\text{MM} = 188.4 \text{ g/mol}$$

2. Consider the reaction of NH_3 and water. The conjugate acid is:

- (A) H_2O
(B) NH_3
(C) CH_3COO^-
(D) H^+
(E) NH_4^+




3. The pH of a buffer system which is 0.450 M HCOOH (aq) and 0.225 M HCOONa (aq) is:

- (A) 3.60×10^{-4}
(B) 4.05
(C) 9.00×10^{-4}
(D) 9.00×10^{-5}
(E) 3.44

$$\text{HCOOH} \rightleftharpoons \text{HCOO}^- + \text{H}^+ \quad K_a = 1.80 \times 10^{-4}$$
$$1.80 \times 10^{-4} = \frac{p}{r} = \frac{[\text{HCOO}^-][\text{H}^+]}{[\text{HCOOH}]}$$
$$1.80 \times 10^{-4} = \frac{(0.225 \text{ M})[\text{H}^+]}{(0.450 \text{ M})}$$
$$[\text{H}^+] = 3.60 \times 10^{-4} \text{ M}$$
$$\text{pH} = -\log[\text{H}^+] = -\log(3.60 \times 10^{-4} \text{ M}) = 3.44$$

4. What is the molar solubility of $\text{Al}(\text{OH})_3$ (aq)? [K_{sp} of $\text{Al}(\text{OH})_3$ is 1.3×10^{-33}]

- (A) $3.6 \times 10^{-12} \text{ M}$
- (B) $2.6 \times 10^{-9} \text{ M}$
- (C) $4.6 \times 10^{-5} \text{ M}$
- (D) $3.6 \times 10^{-17} \text{ M}$
- (E) $2.9 \times 10^{-10} \text{ M}$



$$K_{sp} = 1.3 \times 10^{-33} = [\text{Al}^{3+}] [\text{OH}^{-}]^3 = (x)(3x)^3 = 27x^4$$

$$x = 2.6 \times 10^{-9}$$

Partial credit for (E)
3 points

5. Which of the following is NOT a Lewis base?

- (A) OH^{-}
 - (B) F^{-}
 - (C) $\text{C}_2\text{O}_4^{2-}$
 - (D) Cu^{2+}
 - (E) NH_3
- $\leftarrow \text{Cu}^{2+}$ is a Lewis Acid


6. The concentration of NaOH in water is gradually increased without changing the volume significantly. Which of the following describes what happens to the concentrations of H_3O^{+} and OH^{-} , the pH and the pOH?

	<u>Concentration of H_3O^{+}</u>	<u>Concentration of OH^{-}</u>	<u>pH</u>	<u>pOH</u>
(A)	increases	increases	increases	decreases
(B)	increases	decreases	increases	decreases
(C)	decreases	increases	decreases	increases
(D)	decreases	decreases	decreases	increases
(E)	decreases	increases	increases	decreases

$[\text{OH}^{-}] \uparrow$
 $[\text{H}_3\text{O}^{+}] \downarrow$
 pH \uparrow
 pOH \downarrow

7. What is the coordination number of the metal ion in $[\text{Co}(\text{en})_2(\text{ox})]$?

- (A) 0
- (B) 2
- (C) 4
- (D) 6
- (E) 8



8. The splitting of the d orbitals in a transition metal complex is 171 kJ/mol . What color does this transition metal complex appear to be in transmission?

- (A) green
- (B) blue
- (C) violet
- (D) red
- (E) orange

$$E = \frac{hc}{\lambda} \quad \lambda = \frac{hc}{E} = \frac{(6.626 \times 10^{-34} \text{ J}\cdot\text{s}) (3.00 \times 10^8 \text{ m/s})}{2.84 \times 10^{19} \text{ J/photon}}$$

$$171,000 \frac{\text{J}}{\text{mol}} \left(\frac{1 \text{ mol}}{6.022 \times 10^{23} \text{ photons}} \right) = 2.84 \times 10^{19} \text{ J/photon}$$

$$= 7.00 \times 10^{-7} \text{ m or } 700 \text{ nm}$$

↑
red
↓
appears green

Partial Credit for (D) 3 points

9. Which of the following statements is true about the octahedral complex $[\text{Mn}(\text{CN})_6]^{4-}$?

- (A) The complex is low spin with no unpaired electrons
- (B) The complex is low spin with one unpaired electron
- (C) The complex is low spin with five unpaired electrons
- (D) The complex is high spin with one unpaired electron
- (E) The complex is high spin with five unpaired electrons

$\text{Mn}^{2+} d^5$ ($7-2=5$)
 ↑ strong field
 large Δ_o
 low spin
 — —
 ↑↓ ↑↓ ↑

10. Which of the following processes have a $\Delta S > 0$?

- (A) $\text{CH}_3\text{OH}(l) \rightarrow \text{CH}_3\text{OH}(s)$
- (B) $\text{N}_2(g) + 3 \text{H}_2(g) \rightarrow 2 \text{NH}_3(g)$
- (C) $\text{CH}_4(g) + \text{H}_2\text{O}(g) \rightarrow \text{CO}(g) + 3 \text{H}_2(g)$ 2 moles gas \rightarrow 4 moles gas
- (D) $\text{Na}_2\text{CO}_3(s) + \text{H}_2\text{O}(g) + \text{CO}_2(g) \rightarrow 2 \text{NaHCO}_3(s)$
- (E) All of the above processes have a $\Delta S > 0$.

11. Consider a reaction that has a negative ΔH and a positive ΔS . Which of the following statements is **TRUE**?

- (A) This reaction will be spontaneous only at low temperatures.
- (B) This reaction will be spontaneous at all temperatures.
- (C) This reaction will be nonspontaneous at all temperatures.
- (D) This reaction will be nonspontaneous only at low temperatures.
- (E) All of the above

$$\Delta G = \Delta H - T \Delta S$$

\uparrow (-) (+)

Always (-)

12. Determine the equilibrium constant for a process at 954 K in which $\Delta H_{\text{sys}}^\circ = -90.8 \text{ kJ/mol}$ and $\Delta S_{\text{sys}}^\circ = 102.6 \text{ J/K}$

- (A) 481
- (B) 1.07×10^9
- (C) 2.08×10^{-3}
- (D) 2.14×10^{10}
- (E) 1.94×10^{12}

$$\Delta G = \Delta H - T \Delta S = (-90,800 \text{ J}) - (954 \text{ K})(+102.6 \text{ J/K})$$

$$\Delta G = -188,680 \text{ J}$$



$$\Delta G = -RT \ln K$$

$$-188,680 = (-1)(8.314 \text{ J/mol}\cdot\text{K})(954 \text{ K}) \ln K$$

$$K = 2.14 \times 10^{10}$$

13. When the reaction $\text{Mo (s)} + \text{Cr}_2\text{O}_7^{2-} (\text{aq}) \rightarrow \text{Cr}^{2+} (\text{aq}) + \text{Mo}^{4+} (\text{aq})$ is correctly balanced in acid,

- (A) 4 protons (H^+) are consumed
- (B) 8 protons (H^+) are consumed
- (C) 12 protons (H^+) are consumed
- (D) 14 protons (H^+) are consumed
- (E) 40 protons (H^+) are consumed




14. An electrochemical cell consists of one beaker containing 1.0 M $\text{AgNO}_3(\text{aq})$ into which dips a piece of metallic silver and a second beaker containing 1.0 M $\text{Zn}(\text{NO}_3)_2(\text{aq})$ into which dips a piece of metallic zinc. The beakers are connected by a salt bridge containing 1.0 M $\text{KNO}_3(\text{aq})$. Which of the following statements will be **TRUE** when a wire is connected between the two pieces of metal?

- (A) Electrons will flow from the silver to the zinc. \times
- (B) The cell voltage will be +0.04 V. \times
- (C) There will be no cell voltage as the overall reaction is non-spontaneous. \times
- (D) The silver will plate out on the zinc. \times
- (E) The value of n used to find ΔG° for this cell has the value 2.

$$\Delta G^\circ = -nFE^\circ$$

↑
2



15. A student provides a current of 1.500 amps through an aqueous solution of $\text{Cu}(\text{NO}_3)_2$ for 7.000 hours. The voltage is such that copper is deposited at the cathode. The mass of copper deposited is:



- (A) 0.0804 g
- (B) 0.1608 g
- (C) 6.222 g
- (D) 12.45 g
- (E) 24.89 g

$$7.000 \text{ h} \left(\frac{3600 \text{ s}}{1 \text{ h}} \right) \left(\frac{1.500 \text{ C}}{1 \text{ s}} \right) \left(\frac{1 \text{ mol } e^-}{96,485 \text{ C}} \right) \left(\frac{1 \text{ mol } \text{Cu}^0}{2 \text{ mol } e^-} \right) \left(\frac{63.55 \text{ g}}{1 \text{ mol } \text{Cu}^0} \right) = 12.45 \text{ g Cu}^0$$


↑
Amps

16. A student constructs a voltaic cell from zinc metal and $\text{Zn}(\text{NO}_3)_2$ (aq) and lead metal and $\text{Pb}(\text{NO}_3)_2$ (aq). Which one of the following changes to the cell would cause the cell potential to increase?


- (A) Increase the $\text{Zn}(\text{NO}_3)_2$ (aq) concentration
- (B) Increase the $\text{Pb}(\text{NO}_3)_2$ (aq) concentration
- (C) Increase the mass of $\text{Zn}(s)$
- (D) Decrease the mass of $\text{Zn}(s)$




$$\mathcal{E} = \mathcal{E}^{\circ} - \frac{RT}{nF} \ln \frac{\text{products}}{\text{reactants}}$$



17. Using reduction half-reaction potentials, calculate ΔG° for the following balanced redox reaction:



- (A) $-1.28 \times 10^3 \text{ kJ}$
- (B) -213 kJ
- (C) $-2.21 \times 10^3 \text{ kJ}$
- (D) -2.21 kJ
- (E) -640 kJ



$$\mathcal{E}^{\circ} = (+0.535 \text{ V}) - (-1.676 \text{ V}) = +2.211 \text{ V}$$

$$\Delta G = -nFE^{\circ} = (-1) \times (6 \text{ mol } e^-) \times (96,485 \frac{\text{C}}{\text{mole } e^-}) \times (2.211 \frac{\text{J}}{\text{C}})$$


$$= -1.28 \times 10^6 \text{ J} = -1.28 \times 10^3 \text{ kJ}$$

Partial credit for (B) or (E)
3 points

18. Which of the following statements is **FALSE**?

- (A) Positron emission can be seen as $p \rightarrow n + e^+$
- (B) Electron capture can be seen as $p + e^- \rightarrow n$
- (C) When a beta particle is emitted, an electron is converted to a proton? $e^- \rightarrow ?$
- (D) When an alpha particle is emitted, a helium nucleus is lost from the nucleus of the parent species
- (E) A photon is emitted during gamma emission

19. Consider the figure below. A nuclide in "Region 1" will likely undergo _____ decay and a nuclide in "Region 2" will likely undergo _____ decay.




- (A) alpha; beta
- (B) beta; alpha
- (C) alpha; electron capture
- (D) electron capture; alpha
- (E) beta; positron

Partial Credit
for (B)
3 points


20. Which of the following is NOT an isomer of 2-methyl-3-ethylheptane?

- (A) 5-methylnonane $C_{10}H_{22}$
(B) 2-methyl-4-ethylheptane $C_{10}H_{22}$
(C) 2,2,3,4,4-pentamethylpentane $C_{10}H_{22}$
(D) 2,4-diethyl-3-isopropylpentane $C_{12}H_{26}$
(E) decane $C_{10}H_{22}$




21. An amine and carboxylic acid are reacted together in a condensation reaction. The product is:

- (A) an alcohol
(B) hydrogen peroxide
(C) an amide
(D) an ester
(E) a ketone




22. The addition reaction of 1-hexene and HBr produces:

- (A) 1-hexane
(B) 1-bromohexane
(C) 2-hexane
(D) 1-methylhexane
(E) 2-bromohexane




23. Which of the following lipids will have the lowest melting point?



a)



b)




c)




d)

"most rigid"
 "worst packing"
 Most "double bonds"







24. Xylose, or wood sugar, is one of eight essential sugars for human nutrition. What is the systematic name of this carbohydrate?



Xylose

- (A) ketohexose
- (B) aldohexose
- (C) aldopentose
- (D) ketopentose
- (E) ketotetrose

25. Which of the following is a disaccharide?



monosaccharides linked together


26. Which of the following is **NOT** an example of a secondary protein structure?

- (A) Globular structures such as the four units composing hemoglobin
- (B) α -helix
- (C) β -pleated sheet
- (D) All of the above are secondary protein structures
- (E) None of the above are secondary protein structures


Quaternary (4th Order)

27. Which of the following is responsible for linking two strands of DNA in an α -helix?

- (A) Ester linkages
- (B) Hydrogen bonds
- (C) Glycosidic linkages
- (D) Peptide bonds
- (E) Disulfide linkages




28. The structure of Corticosterone is shown below.




Corticosterone **DOES NOT** contain:

- (A) an alcohol group ~~X~~
- (B) an alkene group ~~X~~
- (C) a ketone group ~~X~~
- (D) an ether group
- (E) a methyl group ~~X~~



29. Which of the following is a secondary alcohol?

- (A) 3-heptanol
- (B) 2-methyl-2-heptanol
- (C) 1-ethanol
- (D) 1-heptanol
- (E) 2,3-dimethyl-1-heptanol



30. Which of the following is **FALSE**?

- (A) A condensation polymer, such as a polyamide, can be formed from a diamine and a dicarboxylic acid
- (B) Polymers are long chains formed by monomer units
- (C) Addition polymerization is common in alkanes *No! alkanes cannot - no pi-system*
- (D) Teflon is a fluorine containing polymer
- (E) Low density polymers are typically more flexible and porous than high density polymers

→ amine - carboxylic acid

31. Amino acids can be linked together with _____ to form _____.

- (A) ester bonds; proteins
- (B) ester bonds; carbohydrates
- (C) ester bonds; polysaccharides
- (D) amide bonds; polysaccharides
- (E) amide bonds; proteins

32. What is the product for the dehydration of $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$?

- (A) $\text{CH}_3\text{CH}_2\text{CH}_2\text{OCH}_2\text{CH}_2\text{CH}_3$
- (B) $\text{CH}_3\text{CH}=\text{CH}_2$
- (C) $\text{CH}_3\text{CH}=\text{CHCH}_3$
- (D) $\text{CH}_3\text{CH}_2\text{CH}=\text{CH}_2$
- (E) $\text{CH}_2=\text{C}=\text{CH}_2$

